

Understanding transversity: present and future

Vincenzo Barone

Università del Piemonte Orientale “A. Avogadro”
and Gruppo Collegato INFN, Alessandria, Italy

Outline

1. What we know about transversity

- Partonic meaning
- QCD evolution
- Non-perturbative results
- k_T -dependent distributions

2. Probing transversity: present and future

- Single-spin asymmetries (SSA): k_T dependence, Collins and Sivers effects, SIDIS and hadroproduction data and their interpretation
- Double-spin asymmetries (DSA): collinear factorization, QCD predictions

3. Summary and conclusions

Transversity in elementary terms

Transversity \equiv Transverse Polarization ($\Pi = \frac{1}{2}\gamma_5\gamma_\perp = \frac{1}{2}\gamma_0\Sigma_\perp$)

Transversity states are linear combinations of helicity states: $|\uparrow\downarrow\rangle = \frac{1}{\sqrt{2}}(|+\rangle \pm i|-\rangle)$

Transversely polarized quarks ($\uparrow\downarrow$) in a transversely polarized proton (\uparrow):


Transversity distribution $\Delta_T f^q(x) = f_\uparrow^q(x) - f_\downarrow^q(x)$ [also called $h_1^q(x)$]

[analogous to the helicity distribution $\Delta f^q(x) = f_+^q(x) - f_-^q(x)$, also called $g_1^q(x)$]

Properties of h_1^q :

- leading twist (unsuppressed)
- chirally odd (non diagonal in helicity basis)
- non-singlet evolution (no gluonic h_1)

Transversity is **not** probed in inclusive DIS: no handbag diagram



To observe transversity, helicity must be flipped twice, so one needs at least **two hadrons** (hadron-hadron scattering or semi-inclusive DIS)

There is **no gluonic transversity distribution** for spin- $\frac{1}{2}$ hadrons: helicity-flip gluon-nucleon amplitudes cannot exist (they would imply $\Delta\lambda = \pm 2$)

A bit of history

Transversity is by now 25 years old, but looks younger (it was forgotten for 10 years)

- Introduced by **Ralston and Soper (1979)** in transversely polarized Drell-Yan
- Rediscovered and studied by **Artru & Mekhfi (1990)**, **Jaffe & Ji (1991)**, **Cortes, Pire & Ralston (1992)**
- See also **Kodaira et al. (1979)**, **Bukhvostov, Kuraev & Lipatov (1981)**, **Efremov & Teryaev (1984)**, **Ratcliffe (1986)**: quark mass contribution to the twist-3 part of g_2
- Theoretical investigation of transversity and related observables: **various authors (1990-2000)**. First experimental results: **E704 (1991)**, **SMC (1999)**, **HERMES (2000)**

[For an overview, see VB, Drago & Ratcliffe, Phys. Rep. 359 (2002) 1]

Highlights of 2002-2004:

- Better understanding of k_T -dependent distributions (Wilson link structures, Sivers function, time-reversal modified universality of parton densities, etc.)
- NLO QCD analyses of transversely polarized collinear hard processes ($p^\uparrow p^\uparrow \rightarrow \gamma X$, $ep^\uparrow \rightarrow e' \Lambda^\uparrow X$, etc.)
- More detailed investigations of transversely polarized lepto- and hadro-production (k_T effects, two-particle final states, higher twists, etc.)
- New processes proposed and studied ($pp^\uparrow \rightarrow j_1 j_2 X$, $pp^\uparrow \rightarrow h_1 h_2 X$, $pp^\uparrow \rightarrow DX$, $p^\uparrow \bar{p}^\uparrow \rightarrow \mu^+ \mu^- (J/\psi) X$, etc.)
- Experiments: data on SSA's in ep^\uparrow (HERMES, COMPASS), new results on SSA in pp^\uparrow at higher energy (STAR)

Notations and terminology

$$\Delta_T f_q(x) \equiv h_1^q(x) \quad \text{transversity distribution function}$$

$$\Delta^N f_{q/p\uparrow}(x, k_T^2) \equiv -\frac{2|\mathbf{k}_T|}{M} f_{1T}^{\perp q}(x, k_T^2) \quad \text{Sivers distribution function}$$


$$\Delta^N D_{h/q\uparrow}(z, P_{hT}^2) \equiv \frac{2|\mathbf{P}_{hT}|}{zM_h} H_1^{\perp q}(z, P_{hT}^2) \quad \text{Collins fragmentation function}$$

Mulders & Tangerman, NP B461 (1996) 197

VB, Drago & Ratcliffe, Phys. Rep. 359 (2002) 1

Bacchetta, D'Alesio, Diehl & Miller, Trento conventions, hep-ph/0410050

The three leading-twist quark distribution functions



number $f_1(x) = \int \frac{d\xi^-}{4\pi} e^{ixP^+\xi^-} \langle P, S | \bar{\psi}(0) \gamma^+ \psi(\xi) | P, S \rangle \Big|_{\xi^+ = \vec{\xi}_\perp = 0}$

helicity $g_1(x) = \int \frac{d\xi^-}{4\pi} e^{ixP^+\xi^-} \langle P, S | \bar{\psi}(0) \gamma^+ \gamma_5 \psi(\xi) | P, S \rangle \Big|_{\xi^+ = \vec{\xi}_\perp = 0}$


transversity $h_1(x) = \int \frac{d\xi^-}{4\pi} e^{ixP^+\xi^-} \langle P, S | \bar{\psi}(0) \gamma^+ \gamma_\perp \gamma_5 \psi(\xi) | P, S \rangle \Big|_{\xi^+ = \vec{\xi}_\perp = 0}$

[Wilson line to ensure gauge invariance: in light-cone gauge, $A^+ = 0$, it reduces to $\mathbb{1}$]

tensor charge: first moment of $h_1 - \bar{h}_1$

$$\langle P, S | \bar{\psi}(0) i\sigma^{\mu\nu} \gamma_5 \psi(0) | P, S \rangle = 2 \delta q S^{[\mu} P^{\nu]}, \quad \delta q = \int_0^1 dx [h_1^q(x) - \bar{h}_1^q(x)]$$

Quark–nucleon helicity amplitudes



$$f_1(x) = f_+(x) + f_-(x) \sim \text{Im} (\mathcal{A}_{++},++ + \mathcal{A}_{+-},+-) = \sum_X (a_{++}^* a_{++} + a_{+-}^* a_{+-})$$

$$g_1(x) = f_+(x) - f_-(x) \sim \text{Im} (\mathcal{A}_{++},++ - \mathcal{A}_{+-},+-) = \sum_X (a_{++}^* a_{++} - a_{+-}^* a_{+-})$$


$$h_1(x) = f_{\uparrow}(x) - f_{\downarrow}(x) \sim \text{Im} \mathcal{A}_{+-},-+ = \sum_X a_{--}^* a_{++}$$

From $\sum_X |a_{++} \pm a_{--}|^2 \geq 0$ follows the **Soffer inequality** [Soffer 1995]:

$$|h_1(x)| \leq \frac{1}{2} [f_1(x) + g_1(x)] = f_+(x)$$

This is an important constraint relating the three leading-twist distributions:

- Must be checked or implemented in modeling h_1



[Chiral quark soliton model, Efremov, Goeke & Schweitzer]

- Provides upper bounds on some transversity observables

QCD evolution of transversity

Purely non-singlet evolution (for spin $\frac{1}{2}$ hadrons): no mixing with gluons

LO (Artru & Mekhfi):

$$\Delta_T \gamma_{qq}^{(0)} = \Delta \gamma_{qq}^{(0)} - \frac{C_F}{n(n+1)}, \quad \Delta_T P_{qq}^{(0)} = \Delta P_{qq}^{(0)} - C_F(1-z)$$


First moment (tensor charge), $\Delta_T \gamma_{qq}^{(0)} = -\frac{2}{3}$:

$$\delta q(Q^2) = \left[\frac{\alpha_S(Q^2)}{\alpha_S(Q_0)^2} \right]^{\frac{4}{27}} \delta q(Q_0^2) \quad \text{decreases slowly}$$

NLO (Hayashigaki, Kanazawa & Koike; Kumano & Miyama; Vogelsang):

$$\delta q(Q^2) = \left[\frac{\alpha_S(Q^2)}{\alpha_S(Q_0)^2} \right]^{\frac{4}{27}} \left[1 - \frac{337}{486 \pi} (\alpha_S(Q_0^2) - \alpha_S(Q^2)) \right] \delta q(Q_0^2)$$

[Hayashigaki, Kanazawa & Koike 1997]



Angular momentum sum rule for transversity [Bakker, Leader & Trueman 2004]:

$$\frac{1}{2} = \frac{1}{2} \sum_{a=q, \bar{q}} \int dx h_1^a(x) + \sum_{a=q, \bar{q}, g} \langle L_T \rangle^a$$

Since transversity **decreases** with increasing Q^2 , the orbital angular momentum must **increase** (assuming an initial zero value) [Ratcliffe]


For spin-1 hadrons, there is also the **gluonic transversity** h_1^g term, which increases with increasing Q^2

It would be extremely interesting to study this sum rule in **pQCD**

Evolution of h_1 rather different from that of g_1 , especially at small x [VB 1997]

$$\text{splitting functions: } \Delta_T P_{qq} \underset{x \rightarrow 0}{\sim} x, \quad \Delta P_{qq} \underset{x \rightarrow 0}{\sim} \text{const.}$$


At small x , h_1 is suppressed compared to g_1



[Hayashigaki, Kanazawa & Koike 1997], $h_1 = g_1$ at the input (GRV) scale

Soffer inequality in QCD: $|h_1(x, Q^2)| \leq \frac{1}{2}[f_1(x, Q^2) + g_1(x, Q^2)]$

- Preserved at LO [VB 1997]
- At NLO, parton distributions are well defined only within a renormalization scheme
- Schemes can be found which preserve the Soffer inequality at NLO [Martin et al. 1998, Bourrely, Soffer and Teryaev 1998]





$R = 2 |h_1| / (f_1 + g_1)$ [Martin, Schäfer, Stratmann & Vogelsang 1998]

Model calculations of h_1 (two examples)


Chiral quark soliton model (Schweitzer et al.)

Chiral chromodielectric model (VB, Calarco & Drago)



- $h_1 \simeq g_1$ at a very low scale (in NRQM, $h_1 = g_1$; in relativistic models the difference is due to lower components of wf's)
- Models differ for the sign of \bar{u} (different procedures to compute antiquark distributions [VB, Calarco, Drago & Simani 2003])

Tensor charges in various models



$$\delta u \sim 0.7 - 1.0, \quad \delta d \sim -(0.1 - 0.4) \quad \text{at } Q^2 = 10 \text{ GeV}^2$$

The relative weight of δu and δd is important for understanding data on single-spin asymmetries in π^\pm, π^0 leptonproduction

Transverse motion of quarks

[Mulders et al., Kotzinian, ...]

k_T -dependent distributions are relevant for single-spin asymmetries

Various possible correlations between \mathbf{k}_T , \mathbf{S}_T and \mathbf{S}_{qT}

Transversely polarized quarks inside a transversely polarized proton:

$$\mathcal{P}_{q\uparrow/p\uparrow}(x, \mathbf{k}_T) = \frac{1}{2} \left\{ f_1(x, \mathbf{k}_T^2) + (\mathbf{S}_T \cdot \mathbf{S}_{qT}) h_1(x, \mathbf{k}_T^2) - \frac{1}{M^2} \left[(\mathbf{k}_T \cdot \mathbf{S}_{qT})(\mathbf{k}_T \cdot \mathbf{S}_T) + \frac{1}{2} \mathbf{k}_T^2 (\mathbf{S}_T \cdot \mathbf{S}_{qT}) \right] h_{1T}^\perp(x, \mathbf{k}_T^2) \right\}$$

$$\mathcal{P}_{q\uparrow/p\uparrow}(x, \mathbf{k}_T) - \mathcal{P}_{q\downarrow/p\uparrow}(x, \mathbf{k}_T) = (\mathbf{S}_T \cdot \mathbf{S}_{qT}) h_1(x, \mathbf{k}_T^2) - \frac{1}{M^2} \left[(\mathbf{k}_T \cdot \mathbf{S}_{qT})(\mathbf{k}_T \cdot \mathbf{S}_T) + \frac{1}{2} \mathbf{k}_T^2 (\mathbf{S}_T \cdot \mathbf{S}_{qT}) \right] h_{1T}^\perp(x, \mathbf{k}_T^2)$$

Both h_1 and h_{1T}^\perp contribute to SSA in ep^\uparrow (via Collins effect), but with different angular distributions, $\sin(\phi_h + \phi_S)$ and $\sin(3\phi_h - \phi_S)$

Unpolarized quarks inside a transversely polarized proton:

$$\mathcal{P}_{q/p\uparrow}(x, \mathbf{k}_T) = f_1(x, \mathbf{k}_T^2) + \frac{(\mathbf{k}_T \times \hat{\mathbf{P}}) \cdot \mathbf{S}_T}{M} f_{1T}^\perp(x, \mathbf{k}_T^2)$$

$$\mathcal{P}_{q/p\uparrow}(x, \mathbf{k}_T) - \mathcal{P}_{q/p\uparrow}(x, -\mathbf{k}_T) = \frac{(\mathbf{k}_T \times \hat{\mathbf{P}}) \cdot \mathbf{S}_T}{M} f_{1T}^\perp(x, \mathbf{k}_T^2)$$

f_{1T}^\perp is the **Sivers distribution function**: azimuthal asymmetry of unpolarized quarks inside a transversely polarized proton

Transversely polarized quarks inside an unpolarized proton:

$$\mathcal{P}_{q\uparrow/p}(x, \mathbf{k}_T) = \frac{1}{2} \left[f_1(x, \mathbf{k}_T^2) + \frac{(\mathbf{k}_T \times \hat{\mathbf{P}}) \cdot \mathbf{S}_{qT}}{M} h_1^\perp(x, \mathbf{k}_T^2) \right]$$

$$\mathcal{P}_{q\uparrow/p}(x, \mathbf{k}_T) - \mathcal{P}_{q\downarrow/p}(x, \mathbf{k}_T) = \frac{(\mathbf{k}_T \times \hat{\mathbf{P}}) \cdot \mathbf{S}_{qT}}{M} h_1^\perp(x, \mathbf{k}_T^2)$$

h_1^\perp is the **Boer-Mulders distribution function**: spin asymmetry of transversely polarized quarks inside an unpolarized proton

“T-odd” distributions f_{1T}^\perp and h_1^\perp

f_{1T}^\perp and h_1^\perp measure T-odd correlations: $(\hat{\mathbf{P}} \times \mathbf{k}_T) \cdot \mathbf{S}_T$ and $(\hat{\mathbf{P}} \times \mathbf{k}_T) \cdot \mathbf{S}_{qT}$


Operator definition of the **Sivers function**:

$$f_{1T}^\perp(x, \mathbf{k}_T^2) \sim \int d\xi^- \int d\xi_T e^{ixP^+ \xi^- - i\mathbf{k}_T \cdot \xi_T} \langle P, S_T | \bar{\psi}(\xi) \gamma^+ W(0, \xi) \psi(0) | P, S_T \rangle$$

If we set the Wilson link W to $\mathbb{1}$, **time reversal** implies [Collins 1993]

$$f_{1T}^\perp(x, \mathbf{k}_T^2) = -f_{1T}^\perp(x, \mathbf{k}_T^2) \quad \Rightarrow \quad \text{Sivers function vanishes}$$

But an explicit calculation [Brodsky, Hwang & Schmidt 2002] shows that




Gluon exchange between struck quark and spectator generates Sivers asymmetry.

Puzzle solved by carefully considering Wilson lines [Collins 2002]

Wilson-line structure of k_T -dependent distributions

$$\mathcal{P}_{q/p}(x, \mathbf{k}_T) = \int \frac{d\xi^-}{4\pi} \int \frac{d\boldsymbol{\xi}_T}{(2\pi)^2} e^{ixP^+\xi^- - i\mathbf{k}_T \cdot \boldsymbol{\xi}_T} \langle P, S | \bar{\psi}(\xi) \gamma^+ W(0, \xi) \psi(0) | P, S \rangle$$

Wilson line $W_\gamma = \text{P exp} \left[-ig \int_\gamma A(x) \cdot dx \right]$: path γ depends on the process



$$\text{SIDIS : } W(0, \xi) = W(0^-, 0_T; \infty^-, 0_T) W(\infty^-, 0_T; \infty^-, \xi_T) W(\infty^-, \xi_T; \xi^-, \xi_T)$$

The gauge potential at ∞^- does not vanish in light-cone gauge [Belitsky, Ji & Yuan 2003]

Time reversal implies

$$\mathcal{P}_{q/p\uparrow}(x, \mathbf{k}_T) \Big|_{\text{future pointing } W} = \mathcal{P}_{q/p\downarrow}(x, \mathbf{k}_T) \Big|_{\text{past pointing } W}$$

SIDIS: Sivers asymmetry from interaction between spectator and **outgoing quark**

DY: Sivers asymmetry from interaction between spectator and **incoming quark**

$$f_{1T}^\perp(x, \mathbf{k}_T^2)_{\text{SIDIS}} = -f_{1T}^\perp(x, \mathbf{k}_T^2)_{\text{DY}}$$

More complicated Wilson link structures in other hard processes studied by [Bomhof, Mulders & Pijlman 2004]. (Time-reversal modified) universality between e^+e^- , DY and SIDIS shown by [Collins & Metz 2004]

At twist 3, effective T-odd distributions emerge from **gluonic poles** [Boer, Mulders & Teryaev 1997]. Connections between \mathbf{k}_T -dependent and twist-3 distributions deserve further study

Probing transversity: taxonomy

To probe transversity one needs at least one extra hadron besides the transversely polarized target

- One hadron in initial state, one in final state (ep scattering):

- $ep^\uparrow \rightarrow e' \pi X$

single spin

- $ep^\uparrow \rightarrow e' \pi^+ \pi^- X$

single spin

- $ep^\uparrow \rightarrow e' \Lambda^\uparrow X$

double spin

- Two hadrons in initial state (pp scattering):

- $p^\uparrow p^\uparrow \rightarrow \mu^+ \mu^- X$

double spin

- $p^\uparrow p^\uparrow \rightarrow \gamma X$

double spin



- $pp^\uparrow \rightarrow h X$

single spin

- Single-spin processes: easier to investigate; asymmetries arise from k_T -dependent distribution and fragmentation functions [except *]
 - $ep^\uparrow \rightarrow e' \pi X$ (Collins and Sivers effects with different angular distributions)
 - $pp^\uparrow \rightarrow h X$ (k_T -factorization only conjectured, not proven; mixing of various asymmetry sources)
 - $ep^\uparrow \rightarrow e' \pi^+ \pi^- X$ [*] (collinear factorization, interference between fragmentation channels)
- Double spin processes: collinear factorization, but experimentally more difficult
 - $ep^\uparrow \rightarrow e' \Lambda^\uparrow X$ (fragmentation into Λ unknown)
 - $p^\uparrow p^\uparrow \rightarrow \mu^+ \mu^- X, p^\uparrow p^\uparrow \rightarrow \gamma X$ (clean, but small asymmetries expected)

Semi-inclusive DIS with a transversely polarized target

$$ep^\uparrow \rightarrow e'hX$$



$$W^{\mu\nu} \sim \text{Tr} [\Phi \gamma^\mu \Xi \gamma^\nu]$$

Ξ fragmentation matrix

Fragmentation matrix (no κ_T): $\Xi(z) \sim D_1(z)\not{P} + \lambda_h G_1(z)\gamma_5\not{P} + H_1(z)\gamma_5\not{S}_T\not{P}$

In theory, the simplest access to transversity is given by the **double-spin process** $ep^\uparrow \rightarrow e' \Lambda^\uparrow X$, which involves

$$H_1(z) = \mathcal{N}_{h^\uparrow/q^\uparrow}(z) - \mathcal{N}_{h^\uparrow/q^\downarrow}(z) \quad [\text{analog of } h_1]$$

Cross-section integrated over \mathbf{P}_T :

$$d\sigma \sim (1-y) \cos(\phi_S + \phi_{S_\Lambda}) h_1(x) H_1(z)$$

Polarization of Λ (measurable by studying the $\Lambda \rightarrow p\pi$ decay):

$$\mathcal{P}_\Lambda = \hat{a}(y) \frac{\sum_q e_q^2 h_1^q(x) H_1^q(z)}{\sum_q e_q^2 f_1^q(x) D_1^q(z)}$$

where $\hat{a}(y) = 2(1-y)/[1+(1-y)^2]$ is the QED depolarizing factor (elementary transverse asymmetry).


The main difficulty in predicting \mathcal{P}_Λ is that $H_{1\Lambda}$ is completely unknown (see some models in [\[Anselmino, Boglione & Murgia 2000\]](#))

In single-spin processes $e p^\uparrow \rightarrow e' \pi^{\pm,0} X$ asymmetries come from transverse motion of quarks (one has to look at \mathbf{P}_{hT} distributions)

Collinear case (λ, η, \dots helicity indices):

$$d\sigma \sim \sum_a \sum_{\lambda\lambda'\eta\eta'} f_a(x) \rho_{\lambda\lambda'} d\hat{\sigma}_{\lambda\lambda'\eta\eta'} \mathcal{D}_{h/a}^{\eta\eta'}(z)$$

partonic subprocess
lepton-quark scattering



Ignoring \mathbf{k}_T and $\boldsymbol{\kappa}_T$, the dependence on the transverse spin of the target is carried by ρ_{+-} , which couples to $\mathcal{D}_{h/a}^{+-}$. Since $\mathcal{D}_{h/a}^{+-}$ vanishes for unpolarized hadrons, there is **no asymmetry in pion production**

Non collinear case (factorization proven by [Ji, Ma & Yuan 2004] for $P_{hT} \ll Q$)

$$d\sigma \sim \sum_a \sum_{\lambda\lambda'\eta\eta'} \mathcal{P}_a(x, \mathbf{k}_T) \rho_{\lambda\lambda'} \otimes d\hat{\sigma}_{\lambda\lambda'\eta\eta'} \otimes \mathcal{D}_{h/a}^{\eta\eta'}(z, \mathbf{\kappa}_T)$$

Single-spin transverse asymmetry are due to

- spin asymmetry of transversely polarized quarks fragmenting into an unpolarized hadron (Collins effect) [non vanishing $\mathcal{D}_{h/a}^{+-}$]
- azimuthal asymmetry of unpolarized quarks inside the transversely polarized proton (Sivers effect) [target transverse spin dependence in ρ_{++}, ρ_{--}]

Fragmentation of transversely polarized quarks into an unpolarized hadron:

$$\mathcal{N}_{h/q\uparrow}(z, \mathbf{P}_{hT}) = D_1(z, \mathbf{P}_{hT}^2) + \frac{(\hat{\mathbf{k}}_T \times \mathbf{P}_{hT}) \cdot \mathbf{S}_{qT}}{zM_h} H_1^\perp(z, \mathbf{P}_{hT}^2)$$

$$\mathcal{N}_{h/q\uparrow}(z, \mathbf{P}_{hT}) - \mathcal{N}_{h/q\downarrow}(z, \mathbf{P}_{hT}) = \frac{(\hat{\mathbf{k}}_T \times \mathbf{P}_{hT}) \cdot \mathbf{S}_{qT}}{zM_h} H_1^\perp(z, \mathbf{P}_{hT}^2)$$

$H_1^\perp(z, \mathbf{P}_{hT}^2)$ is the **Collins fragmentation function**, a T-odd function (not forbidden by time reversal invariance due to final-state interactions)

Distribution of unpolarized quarks inside a transversely polarized proton:

$$\mathcal{P}_{q/p\uparrow}(x, \mathbf{k}_T) - \mathcal{P}_{q/p\uparrow}(x, -\mathbf{k}_T) = \frac{(\mathbf{k}_T \times \hat{\mathbf{P}}) \cdot \mathbf{S}_T}{M} f_{1T}^\perp(x, \mathbf{k}_T^2)$$

$f_{1T}^\perp(x, \mathbf{k}_T^2)$ is the **Sivers distribution function**

Transversely polarized pion leptonproduction cross section:

$$\begin{aligned}
 d\sigma \sim & A(y) \mathcal{I} \left[\frac{\boldsymbol{\kappa}_T \cdot \hat{\mathbf{P}}_{hT}}{M_h} h_1^a H_1^{\perp a} \right] \sin(\phi_h + \phi_S) \\
 & + B(y) \mathcal{I} \left[\frac{\mathbf{k}_T \cdot \hat{\mathbf{P}}_{hT}}{M_h} f_{1T}^{\perp a} D_1^a \right] \sin(\phi_h - \phi_S) \\
 & + C(y) \mathcal{I} \left[\frac{4(\mathbf{k}_T \cdot \hat{\mathbf{P}}_{hT})^2 (\hat{\mathbf{P}}_{hT} \cdot \boldsymbol{\kappa}_T) - 2(\mathbf{k}_T \cdot \hat{\mathbf{P}}_{hT})(\mathbf{k}_T \cdot \boldsymbol{\kappa}_T) - \mathbf{k}_T^2 (\boldsymbol{\kappa}_T \cdot \hat{\mathbf{P}}_{hT})}{2M^2 M_h} h_{1T}^{\perp a} H_1^{\perp a} \right] \\
 & \times \sin(3\phi_h - \phi_S)
 \end{aligned}$$

$$\mathcal{I}[f D] \equiv \int d^2 \mathbf{k}_T \int d^2 \boldsymbol{\kappa}_T \delta^2(\mathbf{k}_T - \mathbf{P}_{hT}/z - \boldsymbol{\kappa}_T) f(x, \mathbf{k}_T^2) D(z, \boldsymbol{\kappa}_T^2)$$

Assuming Gaussian dependence on transverse momenta, $\mathcal{I}[f D] \propto f D$

Variety of angular distributions (two independent physical angles):

- Collins effect: $\sin(\phi_h + \phi_S)$ and [if $\mathbf{k}_T \neq 0$] $\sin(3\phi_h - \phi_S)$
- Sivers effect: $\sin(\phi_h - \phi_S)$

Disentangled by taking azimuthal moments, for instance:

$$\langle \sin(\phi_h + \phi_S) \rangle \equiv \frac{\int d\phi_h d\phi_S \sin(\phi_h + \phi_S) [d\sigma(\phi_h, \phi_S) - d\sigma(\phi_h, \phi_S + \pi)]}{\int d\phi_h d\phi_S [d\sigma(\phi_h, \phi_S) + d\sigma(\phi_h, \phi_S + \pi)]}$$

Other occurrences of transversity (and related functions) in $ep^\uparrow \rightarrow e'\pi X$

Integrated cross-sections at twist 3:

- Unpolarized lepton beam:

$$d\sigma^\uparrow - d\sigma^\downarrow \sim h_1(x) \tilde{H}(z)$$

h_1 couples to the twist-3 fragmentation function \tilde{H}

- Longitudinally polarized lepton beam:

$$d\sigma^{\rightarrow\uparrow} - d\sigma^{\rightarrow\downarrow} \sim h_1(x) \tilde{E}(z)$$

h_1 couples to the twist-3 fragmentation function \tilde{E}

P_{hT} distributions at twist 3:

- Unpolarized lepton beam:

$\sin(2\phi_h - \phi_S)$ asymmetry from coupling of various twist-3 distributions to


Collins function H_1^\perp

HERMES results on single-spin asymmetries in SIDIS

[Airapetian et al. 2004]

$$\langle x \rangle = 0.09, \quad \langle z \rangle = 0.36, \quad \langle Q^2 \rangle = 2.4 \text{ GeV}^2, \quad \langle P_{hT} \rangle = 0.4 \text{ GeV}$$

Collins asymmetries



$A_T^{\pi^+} > 0$, $A_T^{\pi^-} < 0$: consistent with h_1^u positive, h_1^d negative. But $|A_T^{\pi^-}| \gtrsim |A_T^{\pi^+}|$, whereas one expects from models $|h_1^d| \ll |h_1^u|$

Possible explanation: large unfavored Collins functions

$$\pi^+ : \quad 4 h_1^u H_1^{\perp \text{fav}} + h_1^d H_1^{\perp \text{unf}}$$

$$\pi^- : \quad h_1^d H_1^{\perp \text{fav}} + 4 h_1^u H_1^{\perp \text{unf}}$$

π^- data seem to require $H_1^{\perp \text{unf}} \approx -H_1^{\perp \text{fav}}$


Independent information on H_1^{\perp} is very important:

$\Rightarrow e^+e^-$ data from b -factories [Belle]

Other HERMES results on Collins asymmetry:

- Preliminary data on π^0 asymmetry are quite controversial (largely negative $A_T^{\pi^0}$, in conflict with expectations based on isospin invariance)
- $\sin(3\phi_h - \phi_S)$ contribution (due to h_{1T}^{\perp} coupling to Collins function) negligible within errors

Sivers asymmetries



$A_T^{\pi^+} > 0$, $A_T^{\pi^-} \sim 0$: evidence for non vanishing f_{1T}^\perp , but more precise data are needed

COMPASS preliminary results on Collins asymmetry with a deuteron target presented at DIS 04


Kinematics:

$$x \lesssim 0.1 \quad \langle Q^2 \rangle = 2.4 \text{ GeV}^2, \quad \langle P_{hT}^2 \rangle = 0.3 \text{ GeV}^2,$$

The single-spin asymmetry $A_T^{\pi^\pm}$ is found to be compatible with zero (as expected)

Transversity in two-particle leptonproduction

[Collins & Ladinski 1994, Jaffe, Jin & Tang 1998, Bianconi et al. 2000, Bacchetta & Radici 2003]



$$e + p^\uparrow \rightarrow e' + h_1 + h_2 + X$$

$$M_h^2 = (P_1 + P_2)^2, \quad R = \frac{1}{2}(P_1 - P_2)$$

$$z = z_1 + z_2, \quad \xi = z_1/z = 1 - z_2/z$$

Probes angular correlations of the form $(\mathbf{P}_1 \times \mathbf{P}_2) \cdot \mathbf{S}_T = (\mathbf{P}_{hT} \times \mathbf{R}_T) \cdot \mathbf{S}_T$

Transverse polarization of fragmenting quark gives rise to orbital relative motion of h_1 and h_2

Cross section (collinear factorization):

$$d\sigma \sim \sum_q e_q^2 h_1^q(x) I_q(z, M_h^2) \sin(\phi_S + \phi_R)$$

$I_q(z, M_h^2)$ is an **interference fragmentation function**, arising from interference between different production channels


Consider $\pi^+\pi^-$ production via σ, ρ resonance formation [Jaffe, Jin & Tang 1998]:

$$|\pi^+\pi^-\rangle = e^{i\delta_0(M_h)} |\sigma\rangle + e^{i\delta_1(M_h)} |\rho\rangle$$

The interference fragmentation function has the form

$$I(z, M_h^2) \sim \sin \delta_0 \sin \delta_1 \sin(\delta_0 - \delta_1) \hat{I}(z)$$


and changes sign around $M_h = m_\rho$.



In the model of [Radici, Jakob & Bianconi 2002] $I(z, M_h^2)$ has a **Breit-Wigner shape**.

⇒ Preliminary HERMES data

Single-spin asymmetries in hadroproduction



Single transverse asymmetry :
$$A_T = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow}$$

In leading-twist QCD one has $A_T = 0$ (Kane, Pumplin & Repko 1978)

A non-zero asymmetry is generated either by **intrinsic k_T effects**, or by **higher twist effects**

Collinear factorization formula ($\alpha, \gamma \dots$ helicity indices):

$$d\sigma \sim \sum_{abc} \sum_{\alpha\alpha'\gamma\gamma'} \rho_{\alpha'\alpha}^a f_a(x_a) \otimes f_b(x_b) \otimes d\hat{\sigma}_{\alpha\alpha'\gamma\gamma'} \otimes \mathcal{D}_{h/c}^{\gamma'\gamma}(z)$$

For an unpolarized hadron h one has $\mathcal{D}_{h/c}^{\gamma'\gamma} \propto \delta_{\gamma\gamma'}$ and therefore, by helicity conservation, $\alpha = \alpha'$: no dependence on the spin of the target $\rightarrow A_T = 0$

Introduce transverse motion of quarks: **non collinear factorization** (conjectured):

$$d\sigma \sim \sum_{abc} \sum_{\alpha\alpha'\beta\beta'\gamma\gamma'} \rho_{\alpha'\alpha}^a \mathcal{P}_a(x_a, \mathbf{k}_T) \otimes \rho_{\beta'\beta}^b \mathcal{P}_b(x_b, \mathbf{k}'_T) \\ \otimes d\hat{\sigma}_{\alpha\alpha'\beta\beta'\gamma\gamma'} \otimes \mathcal{D}_{h/c}^{\gamma'\gamma}(z, \boldsymbol{\kappa}_T)$$

Three sources of SSA: **Collins effect**, **Sivers effect**, **Boer-Mulders effect**

Collins effect:

$$d\sigma^\uparrow - d\sigma^\downarrow \sim \sum_{abc} [h_1(x_a, \mathbf{k}_T^2) + \frac{\mathbf{k}_T^2}{M^2} h_{1T}^\perp(x_a, \mathbf{k}_T^2)] \otimes f_1(x_b, \mathbf{k}'_T{}^2) \\ \otimes \Delta_{TT} \hat{\sigma}(a^\uparrow b \rightarrow c^\uparrow d) \otimes H_1^\perp(z, \boldsymbol{\kappa}_T^2),$$

In many calculations, only $\boldsymbol{\kappa}_T$ is considered (no h_{1T}^\perp term). A complete treatment of transverse momenta in [Anselmino et al. 2004]

Sivers effect:

$$d\sigma^\uparrow - d\sigma^\downarrow \sim f_{1T}^\perp(x_a, \mathbf{k}_T^2) \otimes f_1(x_b, \mathbf{k}'_T{}^2) \otimes d\hat{\sigma}(ab \rightarrow cd) \otimes D_1(z, \boldsymbol{\kappa}_T^2),$$

Boer-Mulders effect:

$$d\sigma^\uparrow - d\sigma^\downarrow \sim \sum_{abc} [h_1(x_a, \mathbf{k}_T^2) + \frac{\mathbf{k}_T^2}{M^2} h_{1T}^\perp(x_a, \mathbf{k}_T^2)] \otimes h_1^\perp(x_b, \mathbf{k}'_T{}^2) \\ \otimes \Delta_{TT} \hat{\sigma}(a^\uparrow b^\uparrow \rightarrow cd) \otimes D_1(z, \boldsymbol{\kappa}_T^2),$$

The three sources of SSA are entangled in $pp^\uparrow \rightarrow \pi X$ (the only physical angle is $\phi_h - \phi_S$). To disentangle them one would need **less inclusive measurements** (for instance, hadron + jet) [Teryaev].

SSA from higher twists:

$$\begin{aligned} d\sigma^\uparrow - d\sigma^\downarrow \quad \sim \quad & G_F(x_a, y_a) \otimes f_1(x_b) \otimes d\hat{\sigma} \otimes D_1(z) \\ & + h_1(x_a) \otimes E_F(x_b, y_b) \otimes d\hat{\sigma}' \otimes D_1(z) \\ & + h_1(x_a) \otimes f_1(x_b) \otimes d\hat{\sigma}'' \otimes D^{(3)}(z) \end{aligned}$$



$G_F(x, y)$ and $E_F(x, y)$ are quark-gluon correlation functions

$D^{(3)}(z)$ is a twist-3 fragmentation function

E704 results on single-spin asymmetries in $pp^\uparrow \rightarrow \pi X$

[D.L. Adams et al. 1991, Bravar et al. 1996]

$\sqrt{s} = 19.4 \text{ GeV}$, $0.2 \text{ GeV} < \langle P_{\pi T} \rangle < 2.0 \text{ GeV}$



Left: Collins effect with no quark transverse momentum inside the target [Anselmino, Boglione & Murgia 1999]

Right: Collins effect with all transverse momenta taken into account [Anselmino et al. 2004].

The suppression of **Collins effect in pp^\uparrow** is due to **kinematic phases** arising from cross sections of non collinear partonic scattering

$$\sin(\alpha) h_1 H_1^\perp + \sin(\beta) h_{1T}^\perp H_1^\perp$$

Nothing to do with the **magnitude of Collins function**

What about **Collins effect in ep^\uparrow** ?


$$\sin(\phi_h - \phi_S) h_1 H_1^\perp + \sin(3\phi_h - \phi_S) h_{1T}^\perp H_1^\perp$$

In this case there are two **physical angles** and therefore two different angular distributions

STAR results on single-spin asymmetries in $pp^\uparrow \rightarrow \pi^0 X$

[J. Adams et al. 2004]

$$\sqrt{s} = 200 \text{ GeV}, \quad 1.0 \text{ GeV} < \langle P_{\pi T} \rangle < 2.4 \text{ GeV}$$



Well described by: Collins effect, Sivers effect [Anselmino et al.] and higher-twist distribution or fragmentation functions [Qiu & Sterman, Kanazawa & Koike]

E704 vs. (STAR + PHENIX)

- E704 : $\sqrt{s} = 19.4 \text{ GeV}$, $9^\circ < \theta < 67^\circ$
- STAR : $\sqrt{s} = 200 \text{ GeV}$, $\theta = 2.6^\circ$
- PHENIX : $\sqrt{s} = 200 \text{ GeV}$, large $P_{\pi T}$, $\theta \sim 90^\circ$ (central region) $A_T \approx 0$
- [For fixed P_T and \sqrt{s} , large (small) $x_F \Leftrightarrow$ small (large) θ]

- **Unpolarized cross sections:** collinear pQCD correctly reproduce STAR and PHENIX data, but fails to reproduce E704 data (especially at small θ).


Conclusion by [Bourrely & Soffer 2004]: “STAR and E704 asymmetries are different phenomena; E704 asymmetry cannot be attributed to pQCD”

However, **higher twist effects** might be important ($P_{\pi T}$ is not so large)

- [D’Alesio & Murgia 2004]: quark transverse momenta considerably improve agreement with unpolarized cross sections at small \sqrt{s}

Transversity in Drell-Yan processes

$$p^\uparrow + p^\uparrow \rightarrow \mu^+ + \mu^- + X$$



Double transverse asymmetry : $A_{TT}^{DY} = \frac{d\sigma^{\uparrow\uparrow} - d\sigma^{\uparrow\downarrow}}{d\sigma^{\uparrow\uparrow} + d\sigma^{\uparrow\downarrow}}$

At leading order ($q\bar{q}$ annihilation):


$$A_{TT}^{DY} \sim \frac{\sin^2 \theta \cos 2\phi}{1 + \sin^2 \theta} \frac{\sum_q e_q^2 h_1^q(x_1, M^2) \bar{h}_1^q(x_2, M^2) + [1 \leftrightarrow 2]}{\sum_q e_q^2 f_1^q(x_1, M^2) \bar{f}_1^q(x_2, M^2) + [1 \leftrightarrow 2]}$$

Predictions for $A_{TT}^{DY}(pp)$

LO at $\sqrt{s} = 100$ GeV

$h_1 = g_1$ at $Q_0^2 = 0.23$ GeV²


[VB, Calarco & Drago 1997]



NLO at $\sqrt{s} = 40$ GeV (a), 200 GeV (b)

Soffer bound saturated at Q_0

[Martin et al. 1998]



At RHIC energies, very small asymmetries: $A_{TT}^{DY}(pp) \sim 1 - 2\%$

Similar results for prompt photon production: $A_{TT}^{\gamma}(pp) \sim 1\%$ [Mukherjee, Stratmann & Vogelsang 2003]

$$\bar{p}^\uparrow p^\uparrow \rightarrow \mu^+ \mu^- (J/\psi) X \text{ at GSI}$$

$A_{TT}^{DY}(pp)$ at RHIC energies is small because:

1. antiquark transversity distributions are small
2. $\sqrt{s} = 200 \text{ GeV}$, $M < 10 \text{ GeV} \Rightarrow x_1 x_2 = \frac{M^2}{s} < 2.5 \times 10^{-3}$, and at small x transversity evolution is suppressed

[PAX proposal]: polarized antiprotons colliding on polarized protons at HESR of GSI

$$30 \text{ GeV}^2 \lesssim s \lesssim 45 \text{ GeV}^2, \quad M \gtrsim 2 \text{ GeV}, \quad x_1 x_2 \gtrsim 0.1$$


$$A_{TT}^{DY}(\bar{p}p) \sim \frac{\sum_a e_a^2 [h_1^a(x_1, M^2) h_1^a(x_2, M^2) + \bar{h}_1^a(x_1, M^2) \bar{h}_1^a(x_2, M^2)]}{\sum_a e_a^2 [f_1^a(x_1, M^2) f_1^a(x_2, M^2) + \bar{f}_1^a(x_1, M^2) \bar{f}_1^a(x_2, M^2)]}$$

Large asymmetry (**valence-valence combinations**), but small rate

$\Rightarrow J/\psi$ production [Anselmino, VB, Drago, Nikolaev 2004]


- Comparison of J/ψ production in $\bar{p}p$ and pp collisions at $s = 80 \text{ GeV}^2$ shows dominance of $\bar{q}q$ annihilation
- Assume that J/ψ carries over the polarization of the $\bar{c}c$ pair

$$A_{TT}^{J/\psi} \sim \frac{h_1^u(x_1, M_\psi^2) h_1^u(x_2, M_\psi^2)}{f_1^u(x_1, M_\psi^2) f_1^u(x_2, M_\psi^2)}$$



DY at $M = 4 \text{ GeV}$

(solid: $s = 45 \text{ GeV}^2$, dashed: $s = 30 \text{ GeV}^2$)



J/ψ production ($M = 3 \text{ GeV}$)

$A_{TT}^{J/\psi} \sim 0.3$ (similar results by [Efremov, Goeke & Schweitzer 2004])

Summary and conclusions

- Transversity is one of the hottest topics in high-energy spin physics
- From the theoretical point of view, transversity is well understood:
 - QCD evolution known at NLO
 - h_1 and tensor charges sizeable (models, lattice, ...)
 - Related k_T -dependent distributions better known
- Phenomenology of transversity:
 - Single-spin processes are easier to explore but involve unknown functions and require a clear separation of different dynamical mechanisms (Collins, Sivers, higher twists)
 - Double-spin processes are theoretically cleaner and probe transversity more directly, but their experimental investigation is more difficult (it's worth trying!)

- Future perspectives and goals:
 - Theory:
 - * More sophisticated analyses (taking various effects simultaneously into account)
 - * Global studies of different processes (SIDIS, DY, e^+e^- , ...)
 - Experiment:
 - * Extended kinematic ranges (higher P_T and Q , broader x, z, x_F coverage), factorization checks, etc.
 - * Less inclusive measurements (two-particle production, jet+particle production, ...), final state polarimetry (Λ^\uparrow production), antiproton polarization, etc.
- Era of data has arrived. Theory and models on test.

A promising future !